CLINICAL ANATOMY OF THE FEMORAL REGION AND ITS SIGNIFICANCE IN VASCULAR AND SURGICAL PRACTICE

Ashuralieva Gulhumor Kahhorovna
Andijan State Medical Institute, uzbekistan
https://doi.org/10.5281/zenodo.22274772

Abstract

The femoral region is an important anatomical area because it contains major neurovascular structures that connect the pelvis with the lower limb. The femoral artery, femoral vein, femoral nerve, lymphatic vessels, and femoral canal are located within a relatively compact space and have direct clinical significance. The aim of this study was to analyze the clinical anatomy of the femoral region and evaluate its importance in vascular access, surgical procedures, trauma management, lymph-node assessment, and the diagnosis of femoral hernias. A literature-based analytical approach was used to examine the femoral triangle, femoral sheath, vascular and neural relationships, femoral canal, and anatomical landmarks. The analysis demonstrates that the arrangement of the femoral nerve, artery, vein, and lymphatic structures is essential for safe catheterization and surgical intervention. The femoral canal represents a clinically important weak point where femoral hernias may develop, particularly in women. Because femoral hernias have a relatively high risk of incarceration and strangulation, precise anatomical knowledge is necessary for timely diagnosis and treatment. Clinical understanding of the femoral region is therefore essential in surgery, emergency medicine, vascular medicine, radiology, and anatomy education.

Keywords: clinical anatomy, femoral region, femoral triangle, femoral artery, femoral vein, femoral canal, femoral hernia, vascular access.

Introduction

The femoral region is located in the upper anterior part of the thigh and serves as a major passageway for structures traveling between the pelvis and the lower limb.

This area has considerable clinical importance because of the presence of the femoral nerve, artery, vein, lymphatic vessels, and femoral canal.

The femoral triangle is one of the principal anatomical landmarks of the region.

Its boundaries provide an important framework for locating the femoral neurovascular bundle during physical examination and invasive procedures.

The superior boundary of the femoral triangle is formed by the inguinal ligament.

Its lateral boundary is formed by the medial border of the sartorius muscle, while its medial boundary is formed by the medial border of the adductor longus muscle.

Within this region, the femoral nerve, artery, and vein follow a characteristic arrangement.

Knowledge of these relationships is particularly important for arterial puncture, cardiac catheterization, central venous access, vascular surgery, trauma care, and regional anesthesia.

The femoral canal also has important clinical significance because it is a potential site of hernia formation.

The aim of this study was to analyze the major clinical anatomical structures of the femoral region and explain their importance in medical and surgical practice.

Materials and Methods

This study was conducted as a qualitative analytical review of anatomical and clinical literature related to the femoral region.

The analysis focused on the boundaries of the femoral triangle, femoral sheath, femoral artery, femoral vein, femoral nerve, femoral canal, lymphatic structures, and surrounding muscular and fascial anatomy.

Particular attention was given to clinical applications including arterial catheterization, venous access, vascular surgery, femoral nerve block, trauma assessment, lymph-node evaluation, and femoral hernia diagnosis.

The anatomical relationships were interpreted according to their practical relevance in medical procedures and surgical decision-making.

Results

The femoral triangle contains several important structures arranged in a predictable mediolateral pattern.

From lateral to medial, the principal structures are the femoral nerve, femoral artery, femoral vein, and lymphatic structures associated with the femoral canal.

This arrangement is clinically useful when locating the femoral artery pulse or identifying structures during invasive procedures.

The femoral artery is the continuation of the external iliac artery after it passes beneath the inguinal ligament.

It is a major source of blood supply to the lower limb.

The femoral pulse can usually be palpated inferior to the inguinal ligament.

This makes the vessel accessible for diagnostic and therapeutic procedures.

Femoral arterial access is commonly used in angiography, interventional cardiology, and endovascular procedures.

However, puncture must be performed at an appropriate level.

An excessively high puncture may increase the risk of retroperitoneal bleeding, while a very low puncture may involve arterial branches or increase vascular complications.

The femoral vein lies medial to the femoral artery in the proximal part of the femoral triangle.

This relationship is important during femoral venous catheterization.

Because the vein and artery are located close together, accidental arterial puncture can occur.

Ultrasound guidance improves visualization and may increase procedural safety.

The femoral nerve lies lateral to the femoral artery and outside the femoral sheath.

It divides into several muscular and cutaneous branches within the region.

The nerve provides motor innervation to important muscles of the anterior thigh and sensory innervation to parts of the anterior thigh and medial leg.

Injury to the femoral nerve can result in weakness of knee extension and altered sensation.

The femoral sheath is a fascial structure surrounding the proximal portions of the femoral artery and vein.

It is divided into compartments.

The lateral compartment contains the femoral artery, the intermediate compartment contains the femoral vein, and the medial compartment corresponds to the femoral canal.

The femoral canal normally contains loose connective tissue, lymphatic vessels, and a deep inguinal lymph node.

It allows the femoral vein some space for expansion.

However, it is also a potential site for femoral hernia formation.

A femoral hernia occurs when abdominal contents pass through the femoral ring into the femoral canal.

The boundaries of the femoral ring are clinically important.

Anteriorly, it is related to the inguinal ligament.

Posteriorly, it is related to the pectineal ligament and pectineus region.

Medially, it is limited by the lacunar ligament.

Laterally, it is related to the femoral vein.

These rigid boundaries contribute to the risk of incarceration and strangulation when a hernia occurs.

Femoral hernias are generally less common than inguinal hernias but are clinically significant because they can become complicated rapidly.

They are observed more frequently in women, partly because of differences in pelvic anatomy.

The femoral region also contains important lymphatic structures.

Superficial inguinal lymph nodes receive lymph from large areas of the lower limb and external genital region.

Deep inguinal lymph nodes are associated more closely with the femoral vessels.

Lymph-node enlargement in this region may be related to infection, inflammation, or malignant disease.

The femoral region is also important in trauma.

Penetrating or blunt injuries may damage the femoral vessels.

Because these vessels are relatively large, severe hemorrhage can occur rapidly.

Immediate identification and control of vascular injury may therefore be lifesaving.

Discussion

The findings demonstrate that the femoral region has direct relevance to multiple areas of clinical medicine.

Its importance is strongly related to the close organization of large vessels, nerves, and fascial compartments.

The femoral artery is particularly important because of its accessibility.

It provides a convenient route for interventional procedures but also requires precise anatomical localization to minimize complications.

The relationship between the femoral artery and vein is equally important.

Although classical anatomy describes the vein as medial to the artery proximally, their relationship may change distally and may vary among individuals.

This explains why ultrasound guidance has become increasingly useful for vascular procedures.

The femoral nerve is anatomically distinct from the vessels because it lies outside the femoral sheath.

This difference is important during nerve blocks and surgical dissection.

Femoral nerve block can be used for analgesia in selected lower-limb procedures and injuries.

Understanding the location of the nerve relative to the artery improves procedural accuracy.

The anatomy of the femoral canal is especially important in general surgery.

Because the canal is bordered by relatively rigid structures, a hernia sac can become compressed.

This can impair venous drainage and later arterial blood supply to the herniated tissue.

For this reason, femoral hernias may have a greater tendency toward strangulation than many other abdominal wall hernias.

Differentiating a femoral hernia from an inguinal hernia also depends on anatomical landmarks.

Femoral hernias are generally located below the inguinal ligament, while inguinal hernias occur above it.

However, clinical diagnosis may be difficult in obese patients or when the hernia is small.

Ultrasonography and cross-sectional imaging can therefore provide additional information.

The femoral region is also important in emergency medicine.

Major vascular injury can produce rapid blood loss.

Compression of the femoral artery may temporarily reduce distal bleeding while definitive management is prepared.

Clinical anatomy education should therefore connect the topographical organization of the femoral triangle with vascular procedures, trauma, imaging, and hernia surgery.

Conclusion

The femoral region is a clinically significant anatomical area containing major vascular, neural, lymphatic, and fascial structures.

The femoral triangle provides an important landmark for identifying the femoral nerve, artery, vein, and associated structures.

Knowledge of their relationships is essential for arterial and venous catheterization, vascular surgery, regional anesthesia, and trauma management.

The femoral canal is especially important because it is a potential site of femoral hernia formation.

The relatively rigid boundaries of the femoral ring contribute to the risk of incarceration and strangulation.

Accurate understanding of this region can therefore improve diagnostic accuracy and reduce complications during surgical and invasive procedures.

Overall, clinical anatomy of the femoral region demonstrates the importance of integrating structural anatomy with practical medical applications.

References

1. Standring S. *Gray's Anatomy: The Anatomical Basis of Clinical Practice*. 42nd ed. Elsevier; 2020.
2. Moore KL, Dalley AF, Agur AMR. *Clinically Oriented Anatomy*. 9th ed. Wolters Kluwer; 2022.
3. Drake RL, Vogl AW, Mitchell AWM. *Gray's Anatomy for Students*. 5th ed. Elsevier; 2023.
4. Netter FH. *Atlas of Human Anatomy*. 8th ed. Elsevier; 2022.
5. Ellis H, Mahadevan V. *Clinical Anatomy: Applied Anatomy for Students and Junior Doctors*. 14th ed. Wiley-Blackwell; 2019.
6. Tubbs RS, Shoja MM, Loukas M. *Bergman's Comprehensive Encyclopedia of Human Anatomic Variation*. Wiley; 2016.
7. Fitzgibbons RJ Jr, Forse RA. Groin hernias in adults. *New England Journal of Medicine*. 2015;372:756–763.
8. HerniaSurge Group. International guidelines for groin hernia management. *Hernia*. 2018;22:1–165.
9. Brass P, Hellmich M, Kolodziej L, Schick G, Smith AF. Ultrasound guidance versus anatomical landmarks for femoral vein catheterization. *Cochrane Database of Systematic Reviews*. 2015.
10. Snell RS. *Clinical Anatomy by Regions*. 10th ed. Wolters Kluwer; 2019.